

ポスト・ゴンドワナ衝突期に活動した High-K アダカイト：東南極、セール・ロンダーネ山地、ベンゲン花崗岩の例

柚原雅樹¹、亀井淳志²、川野良信³、大和田正明⁴、志村俊昭⁴、東田和弘⁵

¹ 福岡大学

² 島根大学

³ 立正大学

⁴ 山口大学

⁵ 名古屋大学

High-K adakite in post-Gondwana collisional stage: an example of the Vengen Granite, Sør Rondane Mountains, East Antarctica

M. Yuhara¹, A. Kamei², Y. Kawano³, M. Owada⁴, T. Shimura⁴ and K. Tsukada⁵

¹ *Fukuoka University*

² *Shimane University*

³ *Rissho University*

⁴ *Yamaguchi University*

⁵ *Nagoya University*

The Sør Rondane Mountains, located in the eastern Dronning Maud Land, East Antarctica, consists of amphibolite to granulite facies metamorphic rocks and various intrusive rocks (Shiraishi et al., 1997). The early Paleozoic granites are generated during the late- to post-collisional stages as a result of continental collision between the West and East Gondwana would be believed to occur before 620 Ma (Shiraishi et al., 2008; Osanai et al., 2013; Owada et al., 2013). The space-time distribution of igneous activities and their petrogenesis would provide us useful information of geodynamic evolution of a continent-continent collision zone.

The Vengen Granite, one of early Paleozoic granitic rocks, is forming the “Kanino-tume Peak”, the southern end the Vengen ridge of the Sør Rondane Mountains, East Antarctica (Shiraishi et al., 1997). The Vengen Granite is mainly composed of medium-grained mylonitic biotite granite, and intruded into the Main Shear Zone and Kanino-tume Shear Zone. The low-angle cataclasite zone exists in this granitic body. The melamocratic and leucocratic fine-grained two-mica granite dykes are intruded into the Vengen Granite. Part of leucocratic dykes is folded. These dykes have has mylonitic foliation parallel to that of the Vengen Granite. The Vengen Granite is also intruded by quartz veins. Elburg et al. (2016) reported a zircon age of 551 ± 8 Ma by using LA-ICP-MS from this granite.

The Vengen Granite is composed of quartz, K-feldspar, plagioclase, biotite with trace amounts of titanite, allanite, apatite, zircon and opaques as accessory minerals. Parts of rocks contain muscovite. K-feldspar and plagioclase are porphyroclast, up to 12mm in diameter. These porphyroclasts are embedded in smaller biotite, plagioclase, K-feldspar, quartz and accessory minerals. These matrix minerals are up to 2.3mm in diameter. Parts of quartz grains indicate ribbon shape. There are asymmetric pressure shadows in some rocks

SiO₂ contents of the Vengen Granite range from 66.5 to 73.0wt.%. The Vengen Granite indicates much higher K₂O (5.0 - 7.6 wt.%), Ba (1937 - 3288 ppm), Pb (27 - 54 ppm), Rb (142 - 292 ppm), Sr (722 - 1192ppm), Th (24 - 58 ppm), Zr (271 - 448 ppm) contents, and much lower CaO (0.72 - 1.86 wt.%) contents than that of the meta-tonalite (Nils Larsen Tonalite) (Kamei et al., 2013 ; Elburg et al., 2016). The Vengen Granite is plotted on the volcanic arc and syn-collisional granite fields in Rb-(Y+Nb) and Nb-Y tectonic discrimination diagrams (Pearce et al., 1984). The Vengen Granite indicates high Sr/Y ratio (36.1 - 71.5), is plotted on the adakite and Archean high-Al TTGs field in (Sr/Y) -Y diagram. Chondrite-normalized REE patterns of the Vengen Granite are enriched in light REE (LREE) and depleted in heavy REE (HREE). The Vengen Granite do not show Eu anomaly. These patterns similar to that of calc-alkaline meta-tonalite of the Nils Larsen Tonalite corresponding with adakite (Kamei et al., 2013). Model ϵ Sr and ϵ Nd values of the Vengen Granite calculated by 551Ma are 5.6 to 8.6 and -2.3 to -1.1. These values are in that of the tholeiitic meta-tonalite (Kamei et al., 2013).

The Vengen Granite is characterized by high Sr contents and Sr/Y ratios, and corresponds with adakite in Sr/Y-Y diagram. But, this granite indicates high K₂O, Ba, Rb, Zr contents, which are higher than those of common adakite derived from oceanic

slab melting (Martin et al., 2005). The Vengen Granite has high K_2O/Na_2O ratio (1.22 - 1.62 and 3.00) corresponding with adakite derived from lower crust melting (Kamei et al., 2013). Such adakites, which were originated by melting of thickened lower crust, have low MgO, Cr, Co, Ni contents and high Th contents (Xu et al., 2002; Wang et al., 2005; Zhang et al., 2006). Thus, source magma of the Vengen Granite was produced by partial melting in lower parts of thickened continental crust after collision of the West and East Gondwana (Osanai et al., 2013). The source material had isotopic composition similar to the tholeiitic meta-tonalite.

References

- Elburg, M., Andersen, T., Jacobs, J., Läufer, A., Ruppel, A., Krohne, N. and Damaske, D., One Hundred Fifty Million Years of Intrusive Activity in the Sør Rondane Mountains (East Antarctica): Implications for Gondwana Assembly, *Journal of Geology*, 124, 1-26, 2016.
- Kamei, A., Horie, K., Owada, M., Yuhara, M., Nakano, N., Osanai, Y., Adachi, T., Hara, Y., Terao, M., Teuchi, S., Shimura, T., Tsukada, K., Hokada, T., Iwata, C., Shiraishi, K., Ishizuka, H. and Takahashi, Y., Late Proterozoic juvenile arc metatonalite and adakitic intrusions in the Sør Rondane Mountains, eastern Dronning Maud Land, Antarctica. *Precambrian Research*, 234, 47-62, 2013.
- Martin, H., Smithies, R. H., Rapp, R., Moyen, J. -F. and Champion, D., An overview of adakite, tonalite-trondhjemite-granodiorite (TTG), and sanukitoid: relationships and same implications for crustal evolution. *Lithos*, 79, 1-24, 2005.
- Osanai, Y., Nogi, Y., Baba, S., Nakano, N., Adachi, T., Hokada, T., Toyoshima, T., Owada, M., Satish-Kumar, M., Kamei, A. and Kitano, I., Geologic evolution of the Sør Rondane Mountains, east Antarctica: collision tectonics proposed based on metamorphic processes and magnetic anomalies. *Precambrian Research*, 234, 8-29, 2013.
- Owada, M., Kamei, A., Horie, K., Shimura, T., Yuhara, M., Tsukada, K., Osanai Y. and Baba S., Magmatic history and evolution of continental lithosphere of the Sør Rondane Mountains, eastern Dronning Maud Land, East Antarctica. *Precambrian Research*, 234, 63-84, 2013.
- Pearce, J. A., Harris, N. B. W. and Tindle, A. G., Trace element discrimination diagrams for the tectonic interpretation of granitic rocks. *Journal of Petrology*, 25, 956-983, 1984.
- Shiraishi, K., Dunkley, D. J., Hokada, T., Fanning, C. M., Kagami, H. and Hamamoto, T., Geochronological constraints on the Late Proterozoic to Cambrian crust evolution of eastern Dronning Maud Land, East Antarctica: a synthesis of SHRIMP U-Pb age and Nd model age data. In *Geodynamic Evolution of East Antarctica: A Key to the East-West Gondwana Connection* (Satish-Kumar, M., et al., Eds). Geological Society of London, Special Publications, 308, 21-67, 2008.
- Shiraishi, K., Osanai, Y., Ishizuka, H. and Asami, M., Geological map of the Sør Rondane Mountains, Antarctica. Antarctica Geological Map Series, sheet 35, scale 1:250,000, Tokyo, National Institute of Polar Research, 1997.
- Wang, Q., McDermott, F., Xu, J. -F., Bellon, H. and Zhu, Y.-T., Cenozoic K-rich adakitic volcanics in the Hohxil area, northern Tibet: Lower-crustal melting in an intracontinental setting. *Geology*, 33, 465-468, 2005.
- Xu, JF., Shinjo, R., Defant, M. J., Wang, Q. and Rapp, R. P., Origin of Mesozoic adakitic intrusive rocks in the Ningzhen area of east China: Partial melting of delaminated lower continental crust? *Geology*, 30, 1111-1114, 2002.
- Zhang, HF., Zhang, L., Harris, N., Jin, LL. and Yuan, H., U-Pb zircon ages, geochemical and isotopic compositions of granitoids in Songpan-Garze fold belt, eastern Tibetan Plateau: constraints on petrogenesis and tectonic evolution of the basement. *Contributions to Mineralogy and Petrology*, 152, 75-88, 2006.